

## Lecture 1 : Inverse functions

**One-to-one Functions** A function  $f$  is **one-to-one** if it never takes the same value twice or


$$f(x_1) \neq f(x_2) \text{ whenever } x_1 \neq x_2.$$

**Example** The function  $f(x) = x$  is one to one, because if  $x_1 \neq x_2$ , then  $f(x_1) \neq f(x_2)$ .

On the other hand the function  $g(x) = x^2$  is not a one-to-one function, because  $g(-1) = g(1)$ .

**Graph of a one-to-one function** If  $f$  is a one to one function then no two points  $(x_1, y_1)$ ,  $(x_2, y_2)$  have the same  $y$ -value. Therefore no horizontal line cuts the graph of the equation  $y = f(x)$  more than once.

**Example** Compare the graphs of the above functions




### Determining if a function is one-to-one

**Horizontal Line test:** A graph passes the Horizontal line test if each horizontal line cuts the graph at most once.

**Using the graph to determine if  $f$  is one-to-one**

A function  $f$  is one-to-one if and only if the graph  $y = f(x)$  passes the Horizontal Line Test.

**Example** Which of the following functions are one-to-one?



**Using the derivative to determine if  $f$  is one-to-one**

A function whose derivative is always positive or always negative is a one-to-one function. Why?

**Example** Is the function  $g(x) = \sqrt{4x+4}$  a one-to-one function?

## Inverse functions

**Inverse Functions** If  $f$  is a one-to-one function with domain  $A$  and range  $B$ , we can define an inverse function  $f^{-1}$  (with domain  $B$ ) by the rule

$$\boxed{f^{-1}(y) = x \text{ if and only if } f(x) = y.}$$

This is a sound definition of a function, precisely because each value of  $y$  in the domain of  $f^{-1}$  has exactly one  $x$  in  $A$  associated to it by the rule  $y = f(x)$ .

**Example** If  $f(x) = x^3 + 1$ , use the equivalence of equations given above find  $f^{-1}(9)$  and  $f^{-1}(28)$ .

**Note** that the domain of  $f^{-1}$  equals the range of  $f$  and the range of  $f^{-1}$  equals the domain of  $f$ .

**Example** Let  $g(x) = \sqrt{4x + 4}$ . What is Domain  $f$ ?  
What is Range  $g$ ?

Does  $g^{-1}$  exist?

What is Domain  $g^{-1}$ ?

What is Range  $g^{-1}$ ?

What is  $g^{-1}(4)$ ?

### Finding a Formula For $f^{-1}(x)$

Given a formula for  $f(x)$ , we would like to find a formula for  $f^{-1}(x)$ . Using the equivalence

$$x = f^{-1}(y) \text{ if and only if } y = f(x)$$

we can sometimes find a formula for  $f^{-1}$  using the following **method**:

1. In the equation  $y = f(x)$ , if possible solve for  $x$  in terms of  $y$  to get a formula  $x = f^{-1}(y)$ .
2. Switch the roles of  $x$  and  $y$  to get a formula for  $f^{-1}$  of the form  $y = f^{-1}(x)$ .

**Example** Let  $f(x) = \frac{2x+1}{x-3}$ , find a formula for  $f^{-1}(x)$ .

### Composing $f$ and $f^{-1}$ .

We have

$$\text{if } x = f^{-1}(y) \text{ then } y = f(x).$$

Substituting  $f(x)$  for  $y$  in the equation on the left, we get

$$\boxed{f^{-1}(f(x)) = x.}$$

Similarly

$$\text{if } x = f(y) \text{ then } y = f^{-1}(x)$$

and substituting  $f^{-1}(x)$  for  $y$  in the equation on the left, we get

$$\boxed{f(f^{-1}(x)) = x.}$$

**Example** Above, we found that if  $f(x) = \frac{2x+1}{x-3}$ , then  $f^{-1}(x) = \frac{3x+1}{x-2}$ . We can check the above formula for the composition:

$$f(f^{-1}(x)) = f\left(\frac{3x+1}{x-2}\right) = \frac{2\left(\frac{3x+1}{x-2}\right) + 1}{\left(\frac{3x+1}{x-2}\right) - 3} = \frac{(6x+2+x-2)/(x-2)}{(3x+1-3x+6)/(x-2)} = \frac{7x}{7} = x.$$

You should also check that  $f^{-1}(f(x)) = x$ .


### Graph of $y = f^{-1}(x)$

Since the equation  $y = f^{-1}(x)$  is the same as the equation  $x = f(y)$ , the graphs of both equations are identical. To graph the equation  $x = f(y)$ , we note that this equation results from switching the roles of  $x$  and  $y$  in the equation  $y = f(x)$ . This transformation of the equation results in a transformation of the graph amounting to reflection in the line  $y = x$ . Thus

$\boxed{\text{the graph of } y = f^{-1}(x) \text{ is a reflection of the graph of } y = f(x) \text{ in the line } y = x \text{ and vice versa.}}$

**Note** The reflection of the point  $(x_1, y_1)$  in the line  $y = x$  is  $(y_1, x_1)$ . Therefore if the point  $(x_1, y_1)$  is on the graph of  $y = f^{-1}(x)$ , we must have  $(y_1, x_1)$  on the graph of  $y = f(x)$ .

The graphs of  $f(x) = \frac{2x+1}{x-3}$  and  $f^{-1}(x) = \frac{3x+1}{x-2}$  are shown below.



We can derive properties of the graph of  $y = f^{-1}(x)$  from properties of the graph of  $y = f(x)$ , since they are reflections of each other in the line  $y = x$ . For example:

**Theorem** If  $f$  is a one-to-one continuous function defined on an interval, then its inverse  $f^{-1}$  is also one-to-one and continuous. (Thus  $f^{-1}(x)$  has an inverse, which has to be  $f(x)$ , by the equivalence of equations given in the definition of the inverse function.)

**Theorem** If  $f$  is a one-to-one differentiable function with inverse function  $f^{-1}$  and  $f'(f^{-1}(a)) \neq 0$ , then the inverse function is differentiable at  $a$  and


$$(f^{-1})'(a) = \frac{1}{f'(f^{-1}(a))}.$$

**proof**  $y = f^{-1}(x)$  if and only if  $x = f(y)$ . Using implicit differentiation we differentiate  $x = f(y)$  with respect to  $x$  to get

$$1 = f'(y) \frac{dy}{dx} \quad \text{or} \quad \frac{1}{f'(y)} = \frac{dy}{dx}$$

$$\text{or} \quad \frac{1}{f'(y)} = (f^{-1})'(x) \quad \text{or} \quad \frac{1}{f'(f^{-1}(x))} = (f^{-1})'(x)$$

**Geometrically** this means that if  $(a, f^{-1}(a))$  is a point on the curve  $y = f^{-1}(x)$ , then the point  $(f^{-1}(a), a)$  is on the curve  $y = f(x)$  and the slope of the tangent to the curve  $y = f^{-1}(x)$  at  $(a, f^{-1}(a))$  is the reciprocal of the slope of the tangent to the curve  $y = f(x)$  at the point  $(f^{-1}(a), a)$ . The graphs of the function  $f(x) = \frac{2x+1}{x-3}$  and  $f^{-1}(x) = \frac{3x+1}{x-2}$  are shown below. You can verify that  $-7 = (f^{-1})'(3) = \frac{1}{f'(10)}$ .



**Note** To use the above formula for  $(f^{-1})'(a)$ , you do not need the formula for  $f^{-1}(x)$ , you only need the value of  $f^{-1}$  at  $a$  and the value of  $f$  at  $f^{-1}(a)$ .

**Example** Consider the function  $f(x) = \sqrt{4x+4}$  defined above. Find  $(f^{-1})'(4)$ .

What does the formula from the theorem say?

Use the equivalence of the equations  $y = f^{-1}(x)$  and  $x = f(y)$  to find  $f^{-1}(4)$ .

Put this in the formula from the theorem to find  $(f^{-1})'(4)$ .

**Example** Let  $f(x) = x^3 + 1$ , find  $(f^{-1})'(28)$ .

**Example** If  $f$  is a one-to-one function with the following properties:

$$f(10) = 21, \quad f'(10) = 2, \quad f^{-1}(10) = 4.5, \quad f'(4.5) = 3.$$

Find  $(f^{-1})'(10)$ .