

If

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{0}{0}$$

or


$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{\infty}{\infty}$$

Then


$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$$

provided that the latter limit exists.

$$f(x) = f(x_0) + f'(x_0)(x-x_0) + \frac{f''(x_0)}{2!}(x-x_0)^2 + \frac{f'''(x_0)}{3!}(x-x_0)^3 + \frac{f^{(4)}(x_0)}{4!}(x-x_0)^4 + \dots = \sum_{n=0}^{\infty} \frac{f^{(n)}(x_0)}{n!}(x-x_0)^n$$



$$\ln(x) = \int_1^x \frac{1}{t} dt \Rightarrow \ln(2) = \int_1^2 \frac{1}{t} dt \approx 0.69315$$



$$\int u dv = uv - \int v du$$

where it comes from:

The product rule for differentiation

$$\frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}$$

put into reverse

$$\int \frac{d}{dx}(uv) = \int \left(u \frac{dv}{dx} + v \frac{du}{dx} \right)$$


and then

$$uv = \int u \frac{dv}{dx} + \int v \frac{du}{dx}$$

rearranged


$$\int u \frac{dv}{dx} = uv - \int v \frac{du}{dx}$$

Inverse Trigonometry Functions and Their Derivatives




$$\begin{aligned} \frac{d}{dx} \arcsin x &= \frac{1}{\sqrt{1-x^2}} \\ \frac{d}{dx} \arccos x &= \frac{-1}{\sqrt{1-x^2}} \\ \frac{d}{dx} \arctan x &= \frac{1}{1+x^2} \\ \frac{d}{dx} \operatorname{arccot} x &= \frac{-1}{1+x^2} \\ \frac{d}{dx} \operatorname{arcsec} x &= \frac{1}{x\sqrt{x^2-1}} \\ \frac{d}{dx} \operatorname{arccsc} x &= \frac{-1}{x\sqrt{x^2-1}} \end{aligned}$$

The graph of $y = \sin x$ does not pass the horizontal line test, so it has no inverse.



If we restrict the domain (to half a period), then we can talk about an inverse function.



notation: $\arcsin x = \sin^{-1} x = a \sin x$

Definition

$$x = \sin^{-1} y \Leftrightarrow y = \sin x \quad x \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$$

$$x = \cos^{-1} y \Leftrightarrow y = \cos x \quad x \in [0, \pi]$$

notation $\arccos(x) = \cos^{-1} x$

(the angle returned by arcsin fn is between $-\frac{\pi}{2}$ and $\frac{\pi}{2}$)

EX 1 Evaluate these without a calculator.

a) $\cos^{-1}\left(\frac{\sqrt{3}}{2}\right) = \frac{\pi}{6}$



c) $\sin^{-1}(\sin(3\pi/2)) = \sin^{-1}(-1)$




$$= -\frac{\pi}{2}$$

b) $\sin^{-1}(1) = \frac{\pi}{2}$




d) $\cos^{-1}(\cos(-\pi/4)) = \cos^{-1}\left(\frac{\sqrt{2}}{2}\right)$




$$= \frac{\pi}{4}$$

$$y = \tan x$$



$$y = \sec x$$



Definition

$$x = \tan^{-1} y \Leftrightarrow y = \tan x$$

$$x \in (-\pi/2, \pi/2)$$


$$x = \sec^{-1} y \Leftrightarrow y = \sec x$$

$$x \in [0, \pi/2) \cup (\pi/2, \pi]$$

EX 2 Evaluate without a calculator.

a) $\tan^{-1}(-1)$

$= -\frac{\pi}{4}$



A unit circle is drawn on a Cartesian coordinate system. A red arc starts at the point (1, 0) on the positive x-axis and moves clockwise to the point (1/√2, -1/√2) in the fourth quadrant. A blue vector is drawn from the origin to this point. Small 'x' marks are placed at the points (1, 0) and (1/√2, -1/√2).


b) $\sec^{-1}(2)$

$= \cos^{-1}\left(\frac{1}{2}\right)$

$= \frac{\pi}{3}$

$\sec \theta = 2$

$\Leftrightarrow \cos \theta = \frac{1}{2}$




A unit circle is drawn on a Cartesian coordinate system. A red arc starts at the point (1, 0) on the positive x-axis and moves counter-clockwise to the point (1/2, √3/2) in the first quadrant. A blue vector is drawn from the origin to this point. Small 'x' marks are placed at the points (1, 0) and (1/2, √3/2).


c) $\arctan\left(\tan\left(\frac{3\pi}{4}\right)\right)$

$= \arctan(-1)$

$= -\frac{\pi}{4}$




A unit circle is drawn on a Cartesian coordinate system. A red arc starts at the point (1, 0) on the positive x-axis and moves counter-clockwise to the point (-1/√2, 1/√2) in the second quadrant. A blue vector is drawn from the origin to this point. Small 'x' marks are placed at the points (1, 0) and (-1/√2, 1/√2).



Let $\theta = \cos^{-1}x$
 $\Leftrightarrow \cos \theta = x = \frac{x}{1}$
 $\sin(\cos^{-1}x) =$

Pythagorean Thm. $\left\{ \begin{array}{l} \sin \theta \\ = \frac{\sqrt{1-x^2}}{1} \\ = \sqrt{1-x^2} \end{array} \right.$
 $x^2 + d^2 = 1$
 $d^2 = 1 - x^2$
 $d = \sqrt{1-x^2}$
 $(\theta \text{ acute})$



$\theta = \tan^{-1}x$
 $\Leftrightarrow \tan \theta = x = \frac{x}{1}$


$\sec(\tan^{-1}x) =$

$\sec \theta$

$= \frac{\sqrt{1+x^2}}{1}$

$= \sqrt{1+x^2}$

$(\theta \text{ acute})$



$\theta = \sec^{-1}x \Leftrightarrow \sec \theta = x$
 $\cos \theta = \frac{1}{x}$
 $\tan(\sec^{-1}x) =$

$\tan \theta = \frac{\sqrt{x^2-1}}{1} = \sqrt{x^2-1}$

$(\theta \text{ acute})$

EX 3 Calculate $\sin[2\cos^{-1}(1/4)]$ with no calculator.

$$\sin\left(2 \underbrace{\cos^{-1}\left(\frac{1}{4}\right)}_{\theta}\right)$$

$$= \sin(2\theta)$$


$$= 2 \sin\theta \cos\theta$$

(by double angle identity)

$$= 2 \left(\frac{\sqrt{15}}{4}\right) \left(\frac{1}{4}\right) = \boxed{\frac{\sqrt{15}}{8}}$$

$$\text{let } \theta = \cos^{-1}\left(\frac{1}{4}\right)$$

$$\cos\theta = \frac{1}{4}$$



$$?^2 + 1^2 = 4^2$$

$$?^2 = 15$$

$$D_x[\sin x] = \cos x$$

$$D_x[\cos x] = -\sin x$$

$$D_x[\tan x] = \sec^2 x$$

$$D_x[\cot x] = -\csc^2 x$$

$$D_x[\sec x] = \sec x \tan x$$

$$D_x[\csc x] = -\csc x \cot x$$

Derivatives of Inverse Trig Functions

Let $y = \cos^{-1}x$. Find y' .

$$y \in [0, \pi]$$

$$\Leftrightarrow \cos y = x$$

$$D_x(\cos y) = D_x(x)$$

$$-\sin y (y') = 1$$

$$y' = \frac{-1}{\sin y}$$

$$y' = \frac{-1}{\sqrt{1-x^2}}$$

$$y = \cos^{-1}x$$

$$\Rightarrow \sin y = \sin(\cos^{-1}x)$$

$$= \sqrt{1-x^2}$$

$$\Leftrightarrow \boxed{D_x(\cos^{-1}x) = \frac{-1}{\sqrt{1-x^2}}}$$

EX 4 $D_x[\tan^{-1}(5x^2 - 3x + 1)] =$

$$= \frac{1(10x - 3)}{1 + (5x^2 - 3x + 1)^2}$$

$$D_x[\sin^{-1} x] = \frac{1}{\sqrt{1-x^2}}, x \in (-1,1) \quad D_x[\cos^{-1} x] = \frac{-1}{\sqrt{1-x^2}}$$

$$D_x[\sec^{-1} x] = \frac{1}{|x|\sqrt{x^2-1}}, |x| > 1 \quad D_x[\tan^{-1} x] = \frac{1}{1+x^2}$$

EX 5 Evaluate these integrals.

a) $\int_{-1}^1 \frac{1}{1+x^2} dx = \tan^{-1} x \Big|_{-1}^1$
 $= \tan^{-1} 1 - \tan^{-1}(-1)$
 $= \frac{\pi}{4} - (-\frac{\pi}{4}) = \frac{\pi}{2}$

b) $\int \frac{e^x}{1+e^{2x}} dx$

$$\int \frac{1}{\sqrt{1-x^2}} dx = \sin^{-1} x + C$$

$$\int \frac{1}{1+x^2} dx = \tan^{-1} x + C$$

$$\int \frac{1}{x\sqrt{x^2-1}} dx = \sec^{-1} |x| + C$$

$u = e^x \quad \Big| \quad = \int \frac{du}{1+u^2} = \int \frac{1}{1+u^2} du$
 $du = e^x dx$
 note: $e^{2x} = (e^x)^2$

$$= \arctan u + C$$

$$= \boxed{\arctan(e^x) + C}$$

Conclusion

memorize (or write on a reference sheet)

formulas for inverse trig fn derivatives
(* integrals)